

## Anti-Hyperglycemic, Anti-Hyperlipidemic and Antioxidant Potential of Alcoholic-Extract of *Sida cordifolia* (Areal Part) in Streptozotocin-Induced-Diabetes in Wistar-Rats

Mahrukh Ahmad · Shahid Prawez · Mudasir Sultana ·  
Rajinder Raina · Nrip Kishore Pankaj ·  
Pawan Kumar Verma · Shafiqur Rahman

Received: 6 April 2013 / Revised: 21 May 2013 / Accepted: 3 July 2013 / Published online: 14 August 2013  
© The Author(s) 2013. This article is published with open access at Springerlink.com

**Abstract** *Sida cordifolia* is a shrub found throughout the tropical and sub-tropical plains. All parts of the plant are used as anti-rheumatic, antipyretic, anti-asthmatic, laxative, diuretic, vasorelaxative, hypotensive, central nervous system depressant, antioxidant, analgesic and hypoglycemic. The present study was aimed to evaluate the hypoglycemic, anti-hyperlipidemic and antioxidant potential of alcoholic-extract obtained from areal part of *S. cordifolia* in streptozotocin-induced-diabetes in wistar-rats. Diabetes was induced with streptozotocin at the intra-peritoneal dose of 55 mg/kg. Diabetic rats were treated with alcoholic extract of *S. cordifolia* at dosage of 200, 400 mg/kg and glibenclamide (5 mg/kg) after sub-acute administration for 28 days. Alcoholic extract of *S. cordifolia* at 400 mg/kg significantly improved the body-weight whereas significantly decreased the blood glucose level in diabetic rats. However at 400 mg/kg, the alcoholic extract of *S. cordifolia* showed beneficial effect indicating significant decrease in total cholesterol, triglycerides, low density lipids, plasma-creatinine, plasma-urea nitrogen and lipid-peroxidation and a significant increase in high density lipid-level in diabetic rats. Interestingly at 400 mg/kg, a significant increase in antioxidant enzymes such as catalase and superoxide-dismutase-activity was seen in the diabetic

rats. The dose 200 mg/kg of alcoholic extract of *S. cordifolia* showed non-significant change in diabetic rats. The above therapeutic-potential of alcoholic extract of areal parts of plant may be because of the presence of bioactive compounds such as glycosides, resins, alkaloids, sterols, saponins and flavonoids. Thus, the findings of the present study indicate that the alcoholic extract of *S. cordifolia* at dosage level of 400 mg/kg produces anti-diabetic effect in the streptozotocin-induced diabetes in wistar-rats.

**Keywords** *Sida cordifolia* · Alcoholic-extract · Anti-diabetic · Streptozotocin · Wistar-rats

### Introduction

Diabetes mellitus (DM) is the most common endocrine disorder characterized by hyperglycemia resulting from defects either in insulin secretion or insulin action or both [1, 2]. Diabetes is the third killer disease of mankind after cancer and cardiovascular diseases because of its high prevalence, morbidity and mortality [3]. The efficiency of defense mechanism of body is altered in diabetes which results in ineffective scavenging of free radicals and therefore results in tissue damage [4].

To control the disease, several conventional-drugs are available along with insulin but their prolonged use may lead to other complications like blurred vision, hypoglycemia and a lingering condition like coma [5, 6]. The anti-diabetic-drugs such as modern oral hypoglycemic agents' like sulphonyl-ureas (tolbutamide, glibenclamide) and insulin-sensitizer (troglitazone) are associated with various side effects [6]. To reduce its damaging property it is better to use conventional-drugs along with hypoglycemic-herbs [7]. More than 800 plants possess anti-diabetic activity [8,

M. Ahmad · S. Prawez (✉) · M. Sultana · R. Raina ·  
N. K. Pankaj · P. K. Verma  
Division of Pharmacology and Toxicology, FVSc&AH,  
Sher-e-Kashmir University of Agricultural Sciences &  
Technology of Jammu, RS Pura, Jammu 181102, J&K, India  
e-mail: shahidprawez@gmail.com

S. Rahman  
Division of Pathology, FVSc&AH, Sher-e-Kashmir University  
of Agricultural Sciences and Technology of Jammu, RS Pura,  
Jammu 181102, J&K, India

9]. *Sida cordifolia* (Linn.) a shrub belonging to the family Malvaceae, improves the diabetic conditions [10, 11]. It is found through out the tropical and sub tropical plains of India. Its roots, leaves, stem and seeds are used in the folk medicine as anti-rheumatic [12], antipyretic [13], anti-asthmatic [14], laxative, diuretic [12, 15, 16], vaso-relaxative [17], hypotensive [14], CNS depressant [18, 19], antioxidant [20] and hypoglycemic effect [21, 22]. Keeping the fact that diabetes is an emerging problem world-wide and is a major concern of developing countries like India; the present study was conducted with the aim to evaluate the anti-oxidative and anti-diabetic activity of *S. cordifolia* in streptozotocin (STZ) induced diabetes in wistar rats.

## Material and Methods

### Collection and Extraction of Plant Materials

The areal parts of the plant *S. cordifolia* were collected and after authentication plant material was chopped into small pieces and kept under shade for drying (30–35 °C). The plant material was pulverized to powder form with mixer-grinder and subjected to alcoholic extraction in soxhlet-apparatus [23] and its % extractability was determined [24]. The presence of phyto-chemicals in the extract such as alkaloids [25], glycosides [26], saponins [27], sterols [28], resins [29] and flavonoids [30] were determined.

### Experimental Animals

The experimental protocol was duly approved by institutional ethics committee. The study was conducted on healthy wistar rats of either sex weighing 170–230 g, procured from Indian Institute of Integrative Medicine (IIIM Lab), Jammu, India. The animals were provided standard pelleted ration and *ad libitum* drinking tap water. Prior to the start of the experiment, the rats were acclimatized in the laboratory conditions for a period of more than 3 weeks. A daily cycle of 12 h of light and 12 h of darkness was provided to animals.

### Induction of Diabetes

Streptozotocin solution (0.5 %) was freshly prepared in ice cold sodium citrate buffer (0.1 M; pH 4.5) in a volume of 10 mg/ml and administered to overnight fasted wistar-rats intra-peritoneally at the dose of 55 mg kg<sup>-1</sup> body-weight [31–33]. The rats were kept only on glucose solution (5 %) in drinking water for next 24 h after the STZ administration to prevent hypoglycemia [34, 35]. To declare that the rats became diabetic,

after 72 h of STZ administration the biomarker of blood glucose level [36] was determined by using glucometer (Accu-Check, Roche, Germany) and rats showing more than 200 mg/dl blood glucose level were considered as the diabetic rats [37].

### Experimental Design

A total of 30 healthy-wistar rats were selected and divided into five-groups containing six animals in each group. Diabetes was induced in rats of group II, III, IV and V whereas group I served as control. The diabetic rats of group II acted as diabetic control only treated with carboxy-methyl-cellulose (1 %) whereas groups III and IV diabetic rats were treated with alcoholic extract of *S. cordifolia* at dosage of 200 and 400 mg/kg after mixing in carboxy-methyl-cellulose (1 %) for 28 days, respectively. Group V diabetic rats received glibenclamide at dosage of 5 mg/kg orally for 28 days. Blood samples of about 2–4 ml were collected from retro-orbital sinus of wistar-rats under inhalational anesthesia on day 0, 15 and 29 using capillary-tubes and the blood glucose level was measured at the time of sample collection. The blood samples were centrifuged at 3,000 rpm for 15 min to harvest the plasma which was kept in clean sterile glass test tubes at –20 °C for further biochemical analysis. The sediment left after taking out the plasma, from which WBC buffy-coat was removed. The left-over erythrocyte sediment was then washed 2–3 times and diluted with gentle pouring of normal saline solution in the ratio of 1:1 on v/v basis and thoroughly mixed with erythrocyte sediment to make 1 % and 33 % hemolysate used for the estimation of anti-oxidant-enzymes (catalase, superoxide-dismutase) and lipid-peroxidation (LPO) respectively.

### Biochemical and Oxidative Stress Parameters and Body Weight of the Animals

Biochemical indices such as blood glucose, triglycerides, cholesterol, high density lipoproteins (HDL), plasma urea nitrogen and plasma creatinine were assayed on day 0 (pretreatment), day 15 and day 29 (post treatment) using kits (Erba, HP, India). However, low density lipoprotein (LDL) in plasma was estimated using the following formula [38].

$$\text{LDL (mg/dl)} = \text{TC} - \text{HDL} - \text{TG}/5.0,$$

where TC is the total cholesterol and TG is the triglycerides

To detect any changes in body weight the animals were weighed. The antioxidant enzymes superoxide dismutase (SOD) [39], catalase [40] and tissue damage biomarker LPO [41] were estimated in blood samples.

## Statistical Analysis

Statistical data analysis was done using analysis of variance (ANOVA) which was carried out in completely randomized design (CRD) and the significance was tested using Duncan Multiple Range Test [42] and the level of significance was assayed at 5 % level ( $P < 0.05$ ).

## Results and Discussion

The percent alcoholic extractability of *S. cordifolia* was 19.06 % (w/w) and extract revealed the presence of alkaloids, glycosides, saponins, sterols, resins, fixed oil and flavonoids. Kaur et al. [43] also reported similar percent extractability. The findings of phytochemicals were similar to the findings of Ghosal et al. [44], Guntilaka et al. [45], Kapoor [46], Sutradhar et al. [47], and Pawar et al. [48].

A significant increase in blood glucose level was observed in group II, III, IV and V diabetic rats on day 0 as compared to group I control rats (Fig. 1). Treatment was given to group IV diabetic rats with alcoholic extract of *S. cordifolia* at the dose of 400 mg/kg which produced a significant decrease in blood glucose level on day 15 and 29 as compared to day 0. The blood glucose level in group IV and group V on day 29 decreased to the extent which is comparable to control rats of group I, but the dose 200 mg/kg was not enough to decrease blood glucose level appreciably in group III diabetic rats. Although with glibenclamide treatment, the blood glucose level in group V diabetic rats was significantly decreased and was comparable to group I control rats. Similar decrease in blood glucose level was also reported at different doses of *S. cordifolia* [15, 21, 43]. The possible mechanism of hypoglycemic activity of *S. cordifolia* may be through increase in glucose metabolism [43]. It has been reported that mainly alkaloids and flavonoids are responsible for an increase in insulin secretion and peripheral glucose utilization [49].

Diabetes is characterized with the loss of body weight as body protein or fats are being utilized for energy generation through gluconeogenesis [50]. A significant decrease in body weight was found in diabetic rats of group II on day 0 and 29 as compared to rats before diabetic (pretreatment) within same group while in *S. cordifolia* (400 mg/kg) and glibenclamide treated diabetic rats of group IV and V, a significant increase in body weight was found on day 29 as compared to day 0 within the same group, respectively (Fig. 2). Although the dosage of *S. cordifolia* (200 mg/kg) used for diabetic rats of group III was not sufficient to check the decrease in body weight on day 29 as compared to day 0. A significant improvement in body weight of diabetic rats indicated the possible role of extract in


restoration of protein metabolism which is supported by Kaur et al. [43]. The ability of plant extract in restoration of body weight of diabetic rats may be through reversal of gluconeogenesis [51].

A significant decrease in triglyceride (Fig. 3), LDL (Fig. 4) and cholesterol levels (Fig. 5) were found on day 15 and 29 in group IV and V diabetic rats treated with plant extract at dose 400 mg/kg and glibenclamide as compared to day 0 within the group, respectively and also comparable to control rats on day 29. While the dose 200 mg/kg of *S. cordifolia* given to group III diabetic rats showed non significant change in triglyceride and cholesterol level on day 29 as compared to day 0.


A significant increase in HDL level was observed in group IV and group V diabetic rats treated with plant extract at dose 400 mg/kg and glibenclamide on day 15 and 29 as compared to day 0, respectively and comparable to control rats on day 0 (Fig. 6). Although HDL level showed non-significant change in group III diabetic rats treated at 200 mg/kg of plant-extract as compared to day 0. Diabetes affects the fat management indicated by an increase in total cholesterol, triglycerides and LDL, with decrease in HDL levels [52–54]. Similarly, Kaur et al. [43] also reported ameliorative effect with aqueous extract of *S. cordifolia* at the dose of 1,000 mg/kg. The hypo cholesterolemic effect of the plant may be due to overall inhibition of fatty acid synthesis [43, 55]. The significant reduction of LDL levels, in *S. cordifolia* treated rats may be due to the activation of LDL receptors in hepatocytes thus reducing the serum LDL level or may be due to the inhibition of cholesterol synthesis pathway [56]. The effect of *S. cordifolia* extract to decrease triglycerides may be through increase of insulin levels. Insulin activates the enzyme lipoprotein lipase and hydrolyses triglycerides and the deficiency in insulin, thereby causes hyper triglyceridemia [43].

The diabetic hyperglycemia induces elevations of blood creatinine and urea levels which are considered as significant markers of renal dysfunction. A significant decrease in plasma-urea-nitrogen (Fig. 7) and plasma creatinine (Fig. 8) levels was observed on day 29 as compared to day 0 in group IV diabetic rats treated with plant extract at 400 mg/kg and also comparable to control rats of group I. Although the dose 200 mg/kg for group III diabetic rats showed non significant change in plasma urea nitrogen and plasma creatinine levels on day 29. However in group V glibenclamide treated diabetic rats on day 29 a significant decrease in plasma-urea-nitrogen and plasma-creatinine levels was found as compared to day 0 which is comparable to group I control rats. Similar to present findings, Makwana et al. [57] reported that the aqueous extract of *S. cordifolia* has nephro protective potential in nephrotoxicity induced by gentamicin and cisplatin. The higher amount of glucose in blood makes the kidney to work more which


**Fig. 1** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on blood glucose level after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups




**Fig. 2** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on body-weight after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups




**Fig. 3** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on triglyceride after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups




**Fig. 4** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on LDL levels after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups



**Fig. 5** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on cholesterol level after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups




**Fig. 6** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on HDL level after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups






**Fig. 7** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on plasma urea nitrogen after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups



**Fig. 8** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on plasma creatinine after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups




results in more waste product formation indicated by an increase in the serum-creatinine and BUN level [58, 59]. *Sida cordifolia* might have exhibited nephron protective activity by virtue of its antioxidant potential [57].

Oxidative stress in diabetes co-exists with the decrease in antioxidant status, thus increasing the deleterious effects of free radicals. In hyperglycemia, glucose undergoes auto-oxidation which in turn leads to peroxidation of lipids in lipoproteins. A significant decrease in LPO was found on day 29 in group IV and group V diabetes rats treated with plant extract at 400 mg/kg and glibenclamide as compared to day 0 within same group, respectively (Fig. 9). Although group III treated diabetic rats at dose 200 mg/kg of *S. cordifolia* non-significantly decreases the LPO on day 29 as compared to day 0. It is well known that MDA is a terminal product of LPO, thus the concentration of MDA can disclose the extent of LPO in diabetes [60–62]. The reactive oxygen species (ROS) take electrons from polyunsaturated fatty acids of cell membrane which leads to LPO with the loss of cellular functions [63]. Dhalwal et al. [20] reported


that extracts from root of *S. cordifolia* scavenges the free-radical and possess antioxidant-activity. Auddy et al. [64] also reported the *in vivo* and *in vitro* antioxidant activity of whole plant of *S. cordifolia*.

A significant increase in catalase (Fig. 10) and SOD (Fig. 11) activities were found in group IV and group V diabetes-rats treated with plant extract at dose 400 mg/kg and glibenclamide on day 29 as compared to day 0 respectively and enzymes activities were comparable to control rats of group I. Although the dose 200 mg/kg, non-significant changes in antioxidant enzymes (SOD and CAT) were found in group III diabetic rats on day 29 as compared to day 0. Antioxidant enzymes such as SOD and catalase play a very vital role in alleviating the free-radicals oxidative-stress in diabetes. SOD protects from toxic effect of ROS by scavenging superoxide-ions and yielding less reactive hydrogen peroxides [65]. Catalase, a heme protein catalyses the reduction of hydrogen peroxide and protects the tissue from highly reactive hydroxyl radicals [66, 67]. Similar anti-oxidant property of *S. cordifolia* has


**Fig. 9** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on LPO after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups



**Fig. 10** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on antioxidant enzyme catalase activity after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups



**Fig. 11** Effect of alcoholic-extract of *S. cordifolia* and glibenclamide on antioxidant enzyme SOD activity after oral administration in diabetic wistar-rats ( $n = 6$ ). Capital superscript (alphabet) indicates level of significance within the group. Small superscript (alphabet) indicates level of significance between the groups



also been correlated with findings of Pawar et al. [48]. Therefore oxidative stress increases the level of LPO [68] and decrease SOD and catalase levels [60, 61, 69, 70].

*Sida cordifolia* is considered safe as far as its toxicity potential is concerned. As per the literature its LD<sub>50</sub> is more than 3 g/kg [71], thus *S. cordifolia* can be used safely in medicinal practices without causing any toxicity.

## Conclusion

Alcoholic extract of *S. cordifolia* at the dose of 400 mg/kg showed a significant increase in antioxidant enzymes such as catalase and superoxide dismutase levels and significant decrease in LPO in STZ induced diabetes in wistar rats within a treatment period of 28 days. However, no such findings have been found in alcoholic extracts of *S. cordifolia* at dosage level of 200 mg/kg.

Based on the data of present study, it is concluded that alcoholic extract of *S. cordifolia* at a dose of 400 mg/kg have potency to act as anti diabetic, hypoglycemic and anti oxidant properties and also helps to check muscle wasting. Further it also protects from LPO that damages the cell membrane. The above therapeutic potential of the extract from areal part of plants could be due to the presence of bioactive compounds such as glycosides, resins, alkaloids, sterols, saponins, flavanoids etc.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

- David MN (1996) The pathophysiology of diabetic complications. *Ann Intern Med* 124:86–89
- Kumar GPS, Arulselvan P, Kumar SD, Subramaniam PS (2006) Antidiabetic activity of fruits of *Terminalia chebula* on STZ induced diabetic rats. *J Health Sci* 52:283–291
- Gipsen WH, Biessels GJ (2000) Cognition and synaptic plasticity in diabetes mellitus. *Trends Neurosci* 23:542–549
- Nusrath Y, Rajikran E, Sujatha K, Veenavamshee R (2011) Evaluation of renal protective effects of *Desmodium gangeticum* L. in streptozotocin-induced diabetic rats. *Int J Res Pharm Chem* 1:121–127
- Dixit VP, Joshi S (1985) Anti-atherosclerotic effect of alfa-alfa meal injection in chicks: a biochemical evaluation. *Indian J Physiol Pharmacol* 29:47–50
- Shukla R, Sharma SB, Puri D, Prabhu KM, Murthy PS (2000) Medicinal plants for treatment of diabetes mellitus. *Indian J Clin Biochem* 15:169–177
- Ali MR, Akhtar FM (2010) Antihyperglycemic activity of polysaccharides from *Lycium barbarum*. *J Med Plant Res* 4:23
- Alarcon-Aguilara FJ, Roman-Ramos R, Perez-Gutierrez S, Aguilar-Contreras A, Contreras-Weber CCD, Flores-Saenz JL (1998) Study of the anti-hyperglycemic effect of plants used as antidiabetics. *J Ethnopharmacol* 61:101–110
- Cordell GA (2000) Biodiversity and drug discovery: a symbiotic relationship. *Phytochemistry* 55:463–480
- Kirtikar KR, Basu BD (1980) Indian medicinal plants, vol 1. Bishen Singh and Mahendra Pal Singh, Dehradun, p 345
- Fuertes FJ (2006) Antimicrobial activity of the essential oil of *Sida cordifolia* L. *Braz J Pharmacogn* 16:642–644
- Yusuf M, Kabir M (1999) Medicinal plants of Bangladesh. Bangladesh Council of Scientific and Industrial Research, Dhaka, p 226
- Muzzaffer A, Joy Usman A (1991) Screening of *Sida cordifolia* Linn. *Sida rhomboidea* Linn and *Trium fectarotundifolia* Linn. for anti-inflammatory and anti-pyretic activities. *Indian Drugs* 28:397–400
- Medeiros IA, Santos MRV, Nascimento NMS, Duarte JC (2006) Cardiovascular effects of *Sida cordifolia* leaves extract in rats. *Fitoterapia* 77:19–27
- Chopra RN, Handa KL, Kapur LD (1958) Chopra's indigenous drugs of India. Academic Publishers, Meerut, p 409
- Balbach A (1978) A Flora Nacional Na Medicina domestica. MVP Itaquaquecetuba, p 703
- Santos MR, Nascimento NM, Antonioli AR, Medeiros IA (2006) Endothelial derived factors and K<sup>+</sup> channels are involved in the vasorelaxation induced by *Sida cordifolia* leaves in the rat superior mesenteric artery. *Int J Pharm Sci* 61:466–469
- Franco CI, Moraes LC, Quintans-Junior LI, Almeida RN, Antonioli AR (2005) CNS pharmacological effects of hydro-alcoholic extract of *Sida cordifolia* leaves. *J Ethnopharmacol* 98:275–279
- Philip BK, Muralidharan A, Natrajan B, Varadmurthy S, Venkataraman S (2008) Preliminary evaluation of anti-pyretic and anti-ulcerogenic activities of *Sida cordifolia* methanolic extract. *Fitoterapia* 79:229–231
- Dhalwal K, Deshpande YS, Purohit AP, Kadam SS (2005) Evaluation of the antioxidant activity of *Sida cordifolia*. *Pharm Biol* 43:754–761
- Kanth VR, Diwan PV (1999) Analgesic anti-inflammatory and hypoglycemic activity of *Sida cordifolia*. *Phytother Res* 13:75–77
- Franzotti EM, Santos CVF, Rodrigues HMSL, Mourao RHV, Andrade MR, Antonioli AR (2000) Antiinflammatory, analgesic activity and acute toxicity of *Sida cordifolia*. *J Ethnopharmacol* 72:273–277
- Khosla P, Bhanwara S, Singh J, Seth S, Srivastava RK (2000) A study of hypoglycemic effects of *Azadirachta indica* (neem) in normal and alloxan induced diabetic rabbits. *Indian J Physiol Pharmacol* 44:69–74
- Tandle NV (1984) Studies on phytochemical and tranquilising effect of *Ocimum sanctum*. *Indian J Med Res* 14:21–22
- Waldi D (1965) Spray agents for thin-layer chromatography. In: Stahl E (ed) Thin-layer chromatography. Springer, Berlin, p 496
- Ramakrishnan S, Prasanna KG, Rajan R (1994) Text book of medical biochemistry, 4th edn. Orient Longman, New Delhi, p 582
- Kokate CK, Purohit AP, Gokhale SB (1999) Pharmacognosy. Nirali Prakashan, Pune, p 549
- Trease GE, Evans WC (2002) Pharmacognosy, 15th edn. Saunders Publishers, London, p 42
- Sofowora A (1982) Medicinal plants and traditional medicine in Africa, 1st edn. Wiley, Chichester, p 256
- Mace ME (1963) Histochemical localisation of phenols in healthy and diseased banana roots. *Physiol Plantarum* 16:915–925
- Chang KJ (2000) Effect of taurine and beta alanine on morphological changes of pancreas in streptozotocin induced diabetic rats. *Adv Exp Med Biol* 483:571–577



32. Ramesh B, Pugalendi KV (2006) Impact of umbelliferone (7-hydroxycoumarin) on hepatic marker enzymes in STZ diabetic rats. *Indian J Pharmacol* 38:209–210
33. Gayathri M, Kannabiran K (2008) Hypoglycemic activity of *Hemidesmus indicus* on STZ induced diabetic rats. *Int J Diabetes Dev Ctries* 28:6–10
34. Akbarzadeh A, Norouzzian D, Mehrabi MR, Jamshidi S, Farhangi A, Allahverdi A, Mofidian SMA, Lame BR (2007) Induction of diabetes by STZ in rats. *Indian J Clin Biochem* 22:60–64
35. Abeeleh AM, Ismail ZB, Alzaben KR, Abu-Halaweh SA, Al-Essa MK, Abuabeeleh J, Alsmady MM (2009) Induction of diabetes mellitus in rats using intraperitoneal STZ. *Eur J Sci Res* 32:398
36. Karunanayake EH, Hearse DJ, Mellows G (1975) The metabolic fate and elimination of STZ. *Biochem Soc Trans* 3:410–414
37. Sokeng SD, Lontsi D, Moundipa PF, Jatsa HB, Watcho P, Kamthoung P (2007) Hypoglycemic effect of *Anacardium occidentale* L. methanol extract and fractions on streptozotocin-induced diabetic rats. *Glob J Pharmacol* 1:01–05
38. Friedewald WT, Levy RI, Fredrickson DS (1972) Estimation of the concentration of low density lipoproteins in plasma without the use of preparative ultracentrifuge. *Clin Chem* 18:449–452
39. Marklund S, Marklund M (1974) Involvement of superoxide anion radicle in autoxidation of pyrogallol and a convenient assay of superoxide dismutase. *Eur J Biochem* 47:469–474
40. Aebi HE (1984) Catalase in vitro. *Methods Enzymol* 105:121–126
41. Shafiq-ur-Rehman (1984) Lead induced regional lipid peroxidation in brain. *Toxicol Lett* 21:333–337
42. Duncan DB (1955) Multiple range and multiple F test. *Biometrics* 11:1–42
43. Kaur G, Kamboj P, Kalia AN (2011) Antidiabetic and anti-hypercholesterolemic effects of aerial parts of *Sida cordifolia* Linn. on STZ-induced diabetic rats. *Indian J Nat Prod Res* 2:428–434
44. Ghosal S, Chauhan RRPS, Mehta R (1975) Alkaloids of *Sida cordifolia*. *Phytother Chem* 14:830–832
45. Gunatilaka AAL, Sotheeswaran S, Balasubramanian S, Chandrasekara AI, Sriyani HTB (1980) Studies on medicinal plants of Sri Lanka. *Planta Med* 39:66–72
46. Kapoor LD (1990) Handbook of ayurvedic medicinal plant, vol 303. CRC, Boca Raton, p 26
47. Sutradhar RK, Rahman AKMM, Ahmad M, Bachar SC, Saha A, Guha SK (2006) Analgesic and anti-inflammatory principles from *Sida cordifolia* L. *J Biol Sci* 6:160–163
48. Pawar RS, Jain A, Sharma P, Chaurasiya PK, Singour PK (2011) In vitro studies on *Sida cordifolia* Linn for anthelmintic and antioxidant properties. *Chin Med* 2:47–52
49. Ram KK, Suresh K, Sunil S (2012) Evaluation of analgesic activity of various extracts of *Sida tiagii* bhandari. *Acta Pol Pharm* 69:1103–1109
50. Jayasri MA, Radha A, Mathew TL (2009)  $\alpha$ -Amylase and  $\alpha$ -glucosidase inhibitory activity of *Costus pictus* D. DON in the management of diabetes. *J Herb Med Toxicol* 3:91–94
51. Sharma B, Santosh K, Satapathi A, Roy P (2007) Hypoglycemic and hypolipidemic effect of *Aegle marmelos* (L) leaf extract on STZ induced diabetic mice. *Indian J Pharmacol* 3:444–452
52. Babu PS, Srinivasan K (1997) Hypolipidemic action of curcumin, the active principle of *Curcuma longa* in STZ induced diabetic rats. *Biomed Life Sci* 166:169–175
53. Guerci B, Antebi H, Meyer L, Durlach V, Ziegler O, Nicolas J, Alcindor L, Drouin P (1999) Increased ability of LDL from normolipidemic type-II diabetic women to generate peroxides. *Clin Chem* 45:1439
54. Arvind K, Pradeepa R, Deepa R, Mohan V (2002) Diabetes and coronary artery disease. *Indian J Med Res* 116:163–176
55. Chi MS, Koh ET, Stewart TJ (1982) Effect of garlic on lipid metabolism of rats fed with cholesterol or lard. *J Nat* 112:241–248
56. Rang HP, Dale MM, Ritter JM (1999) Textbook of pharmacology. Churchill Livingstone, Edinburgh, pp 301–305
57. Makwana VM, Pandya NM, Darji ND, Desai AS, Bhaskar VH (2012) Assessment of nephroprotective potential of *Sida cordifolia* L. in experimental animals. *Der Pharm Lett* 4:175–180
58. Khan AM, Sultana M, Raina R, Dubey N, Verma PK (2013) Effect of sub-acute oral exposure of Bifenthrin on biochemical parameters in crossbred goats. *Proc Natl Acad Sci India Sect B Biol Sci*. doi:10.1007/s40011-012-0150-x
59. Home P, Mant J, Diaz Z, Turner C (2008) Management of type 2 diabetes: summary of updated NICE guidance. *Br Med J* 336:1306–1308
60. Baynes JW (1991) Role of oxidative stress in development of complications of diabetes mellitus. *Diabetes* 40:405–412
61. Kakkar R, Kalra J, Mantha SV, Prasad K (1995) Lipid peroxidation and activity of antioxidant enzymes in diabetic rats. *Mol Cell Biochem* 151:113–119
62. Mohammed AK, Bierhaun A, Schiekofer S, Tritschler H, Ziegler H, Nawroth PP (1999) The role of oxidative stress and NF (B) activation in late diabetic complications. *Biofactors* 10:175–179
63. Maxwell SR (1995) Prospects for the use of antioxidant therapies. *Drugs* 49:345–361
64. Auddy B, Ferreira M, Blasina F, Lafon L, Arredondo F, Dajas F, Tripathi PC, Seal T, Mukherjee B (2003) Screening of antioxidant activity of three Indian medicinal plants, traditionally used for the management of neurodegenerative diseases. *J Ethnopharmacol* 84:131–138
65. Zakaryan AE, Aivazyazyan NM, Karagezyan KG (2002) Comparative analysis of the superoxide dismutase activity in tissues of higher vertebrates. *Dokl Biochem Biophys* 382:13–15
66. Chelikani P, Fita I, Loewen PC (2004) Diversity of structures and properties among catalase. *Cell Mol Life Sci* 61:192–208
67. Mc-cord JM, Keele BB, Fridovich I (1976) An enzyme based theory of obligate anaerobiosis, the physiological functions of superoxide dismutase. *Proc Nat Acad Sci USA* 68:1024–1027
68. Atalay M, Laaksonen DE (2002) Diabetes, oxidative stress and physical exercise. *Indian J Sports Sci Med* 1:1–4
69. Maritim AC, Sanders RA, Watkin JB (2003) Diabetes, oxidative stress, antioxidants: a review. *J Biochem Mol Toxicol* 17:24–38
70. Moussa SA (2008) Oxidative stress in diabetes mellitus. *Rom J Biophys* 18:225–230
71. K Konali, Bassoli INH, Hilou A, Aworet-samseny RRR, Souza A, Nicolas B, Dicko MH, Datte JY, Batchi BM (2012) Toxicity assessment and analgesic activity investigation of aqueous acetone extracts of *Sida acuta* Burn f. and *Sida cordifolia* L (Malvaceae) and medicinal plants of Burkina Faso. *BMC Complem Altern Med* 12:120